Cambridge Assessment International Education

Cambridge International AS & A Level

CANDIDATE NAME									
CENTRE NUMBER						CANDIDATE NUMBER			

1939478240

MATHEMATICS **9709/12**

Paper 1 Pure Mathematics 1 **February/March 2020**

1 hour 50 minutes

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [ ].

This document has 4 pages.

© UCLES 2020


1 The function f is defined by $\mathrm{f}(x) = \dfrac{1}{3x+2} + x^2$ for $x < -1$.

Determine whether f is an increasing function, a decreasing function or neither. [3]

2 The graph of $y = \mathrm{f}(x)$ is transformed to the graph of $y = 1 + \mathrm{f}\left(\frac{1}{2}x\right)$.

Describe fully the two single transformations which have been combined to give the resulting transformation. [4]

3



The diagram shows part of the curve with equation $y = x^2 + 1$. The shaded region enclosed by the curve, the y-axis and the line $y = 5$ is rotated through 360° about the ***y*-axis**.

Find the volume obtained. [4]

4 A curve has equation $y = x^2 - 2x - 3$. A point is moving along the curve in such a way that at P the y-coordinate is increasing at 4 units per second and the x-coordinate is increasing at 6 units per second.

Find the x-coordinate of P. [4]

5 Solve the equation

$$\frac{\tan\theta + 3\sin\theta + 2}{\tan\theta - 3\sin\theta + 1} = 2$$


for $0° \leqslant \theta \leqslant 90°$. [5]

6 The coefficient of $\dfrac{1}{x}$ in the expansion of $\left(2x + \dfrac{a}{x^2}\right)^5$ is 720.

(a) Find the possible values of the constant a. [3]

(b) Hence find the coefficient of $\frac{1}{x^7}$ in the expansion. [2]

7



The diagram shows a sector AOB which is part of a circle with centre O and radius 6 cm and with angle $AOB = 0.8$ radians. The point C on OB is such that AC is perpendicular to OB. The arc CD is part of a circle with centre O, where D lies on OA.

Find the area of the shaded region. [6]

8 A woman's basic salary for her first year with a particular company is \$30 000 and at the end of the year she also gets a bonus of \$600.

(a) For her first year, express her bonus as a percentage of her basic salary. [1]

At the end of each complete year, the woman's basic salary will increase by 3% and her bonus will increase by \$100.

(b) Express the bonus she will be paid at the end of her 24th year as a percentage of the basic salary paid during that year. [5]

9 **(a)** Express $2x^2 + 12x + 11$ in the form $2(x + a)^2 + b$, where a and b are constants. [2]

The function f is defined by $\mathrm{f}(x) = 2x^2 + 12x + 11$ for $x \leqslant -4$.

(b) Find an expression for $\mathrm{f}^{-1}(x)$ and state the domain of f^{-1}. [3]

The function g is defined by $\mathrm{g}(x) = 2x - 3$ for $x \leqslant k$.

(c) For the case where $k = -1$, solve the equation $\mathrm{fg}(x) = 193$. [2]

(d) State the largest value of k possible for the composition fg to be defined. [1]

© UCLES 2020

10 The gradient of a curve at the point (x, y) is given by $\frac{dy}{dx} = 2(x+3)^{\frac{1}{2}} - x$. The curve has a stationary point at $(a, 14)$, where a is a positive constant.

(a) Find the value of a. [3]

(b) Determine the nature of the stationary point. [3]

(c) Find the equation of the curve. [4]


11 **(a)** Solve the equation $3\tan^2 x - 5\tan x - 2 = 0$ for $0° \leqslant x \leqslant 180°$. [4]

(b) Find the set of values of k for which the equation $3\tan^2 x - 5\tan x + k = 0$ has no solutions. [2]

(c) For the equation $3\tan^2 x - 5\tan x + k = 0$, state the value of k for which there are three solutions in the interval $0° \leqslant x \leqslant 180°$, and find these solutions. [3]

12 A diameter of a circle C_1 has end-points at $(-3, -5)$ and $(7, 3)$.

(a) Find an equation of the circle C_1. [3]



The circle C_1 is translated by $\begin{pmatrix} 8 \\ 4 \end{pmatrix}$ to give circle C_2, as shown in the diagram.

(b) Find an equation of the circle C_2. [2]

The two circles intersect at points R and S.

(c) Show that the equation of the line RS is $y = -2x + 13$. [4]

(d) Hence show that the x-coordinates of R and S satisfy the equation $5x^2 - 60x + 159 = 0$. [2]

© UCLES 2020