Cambridge Assessment International Education

Cambridge International AS & A Level

CANDIDATE NAME									
CENTRE NUMBER						CANDIDATE NUMBER			

1862605169

MATHEMATICS **9709/33**

Paper 3 Pure Mathematics 3 **May/June 2020**

1 hour 50 minutes

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [ ].

This document has 4 pages.

© UCLES 2020

1 Solve the inequality $|2x - 1| > 3|x + 2|$. [4]

2 Find the exact value of $\int_0^1 (2 - x)e^{-2x}\,dx$. [5]

3 **(a)** Show that the equation

$$\ln(1 + e^{-x}) + 2x = 0$$

can be expressed as a quadratic equation in e^x. [2]

(b) Hence solve the equation $\ln(1 + e^{-x}) + 2x = 0$, giving your answer correct to 3 decimal places. [4]

4 The equation of a curve is $y = x\tan^{-1}\left(\frac{1}{2}x\right)$.

(a) Find $\frac{dy}{dx}$. [3]

(b) The tangent to the curve at the point where $x = 2$ meets the y-axis at the point with coordinates $(0, p)$.

Find p. [3]

5 By first expressing the equation

$$\tan\theta\tan(\theta + 45°) = 2\cot 2\theta$$

as a quadratic equation in $\tan\theta$, solve the equation for $0° < \theta < 90°$. [6]

6 **(a)** By sketching a suitable pair of graphs, show that the equation $x^5 = 2 + x$ has exactly one real root. [2]

(b) Show that if a sequence of values given by the iterative formula

$$x_{n+1} = \frac{4x_n^5 + 2}{5x_n^4 - 1}$$

converges, then it converges to the root of the equation in part **(a)**. [2]

(c) Use the iterative formula with initial value $x_1 = 1.5$ to calculate the root correct to 3 decimal places. Give the result of each iteration to 5 decimal places. [3]

7 Let $\mathrm{f}(x) = \dfrac{2}{(2x-1)(2x+1)}$.

(a) Express $\mathrm{f}(x)$ in partial fractions. [2]

(b) Using your answer to part **(a)**, show that

$$\big(\mathrm{f}(x)\big)^2 = \frac{1}{(2x-1)^2} - \frac{1}{2x-1} + \frac{1}{2x+1} + \frac{1}{(2x+1)^2}.$$ [2]

(c) Hence show that $\displaystyle\int_1^2 \big(\mathrm{f}(x)\big)^2 \,\mathrm{d}x = \tfrac{2}{5} + \tfrac{1}{2}\ln\left(\tfrac{5}{9}\right)$. [5]

8 Relative to the origin O, the points A, B and D have position vectors given by

$$\overrightarrow{OA} = \mathbf{i} + 2\mathbf{j} + \mathbf{k}, \quad \overrightarrow{OB} = 2\mathbf{i} + 5\mathbf{j} + 3\mathbf{k} \quad \text{and} \quad \overrightarrow{OD} = 3\mathbf{i} + 2\mathbf{k}.$$

A fourth point C is such that $ABCD$ is a parallelogram.

(a) Find the position vector of C and verify that the parallelogram is not a rhombus. [5]

(b) Find angle BAD, giving your answer in degrees. [3]

(c) Find the area of the parallelogram correct to 3 significant figures. [2]

9 **(a)** The complex numbers u and w are such that

$$u - w = 2\mathrm{i} \quad \text{and} \quad uw = 6.$$


Find u and w, giving your answers in the form $x + \mathrm{i}y$, where x and y are real and exact. [5]

(b) On a sketch of an Argand diagram, shade the region whose points represent complex numbers z satisfying the inequalities

$$|z - 2 - 2\mathrm{i}| \leqslant 2, \quad 0 \leqslant \arg z \leqslant \tfrac{1}{4}\pi \quad \text{and} \quad \mathrm{Re}\, z \leqslant 3.$$ [5]

© UCLES 2020

10



A tank containing water is in the form of a hemisphere. The axis is vertical, the lowest point is A and the radius is r, as shown in the diagram. The depth of water at time t is h. At time $t = 0$ the tank is full and the depth of the water is r. At this instant a tap at A is opened and water begins to flow out at a rate proportional to $\sqrt{h}$. The tank becomes empty at time $t = 14$.

The volume of water in the tank is V when the depth is h. It is given that $V = \frac{1}{3}\pi(3rh^2 - h^3)$.

(a) Show that h and t satisfy a differential equation of the form

$$\frac{\mathrm{d}h}{\mathrm{d}t} = -\frac{B}{2rh^{\frac{1}{2}} - h^{\frac{3}{2}}},$$

where B is a positive constant. [4]

(b) Solve the differential equation and obtain an expression for t in terms of h and r. [8]

© UCLES 2020