Cambridge Assessment International Education

Cambridge International AS & A Level

CANDIDATE NAME									
CENTRE NUMBER						CANDIDATE NUMBER				

3602942828

MATHEMATICS **9709/31**

Paper 3 Pure Mathematics 3 **May/June 2021**

1 hour 50 minutes

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [ ].

This document has 3 pages.

© UCLES 2021

[Turn over

PapaCambridge

1 Solve the inequality $2|3x - 1| < |x + 1|$. [4]

2 Find the real root of the equation $\dfrac{2e^x + e^{-x}}{2 + e^x} = 3$, giving your answer correct to 3 decimal places. Your working should show clearly that the equation has only one real root. [5]

3 **(a)** Given that $\cos(x - 30°) = 2\sin(x + 30°)$, show that $\tan x = \dfrac{2 - \sqrt{3}}{1 - 2\sqrt{3}}$. [4]

(b) Hence solve the equation

$$\cos(x - 30°) = 2\sin(x + 30°),$$

for $0° < x < 360°$. [2]

4 **(a)** Prove that $\dfrac{1 - \cos 2\theta}{1 + \cos 2\theta} \equiv \tan^2 \theta$. [2]

(b) Hence find the exact value of $\displaystyle\int_{\frac{1}{6}\pi}^{\frac{1}{3}\pi} \frac{1 - \cos 2\theta}{1 + \cos 2\theta}\, d\theta$. [4]

5 **(a)** Solve the equation $z^2 - 2piz - q = 0$, where p and q are real constants. [2]

In an Argand diagram with origin O, the roots of this equation are represented by the distinct points A and B.

(b) Given that A and B lie on the imaginary axis, find a relation between p and q. [2]

(c) Given instead that triangle OAB is equilateral, express q in terms of p. [3]


6 The parametric equations of a curve are

$$x = \ln(2 + 3t), \qquad y = \frac{t}{2 + 3t}.$$

(a) Show that the gradient of the curve is always positive. [5]

(b) Find the equation of the tangent to the curve at the point where it intersects the y-axis. [3]

7



The diagram shows the curve $y = \dfrac{\tan^{-1} x}{\sqrt{x}}$ and its maximum point M where $x = a$.

(a) Show that a satisfies the equation

$$a = \tan\left(\frac{2a}{1 + a^2}\right).$$ [4]

(b) Verify by calculation that a lies between 1.3 and 1.5. [2]

(c) Use an iterative formula based on the equation in part **(a)** to determine a correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

8 With respect to the origin O, the points A and B have position vectors given by $\overrightarrow{OA} = \begin{pmatrix} 1 \\ 2 \\ 1 \end{pmatrix}$ and $\overrightarrow{OB} = \begin{pmatrix} 3 \\ 1 \\ -2 \end{pmatrix}$. The line l has equation $\mathbf{r} = \begin{pmatrix} 2 \\ 3 \\ 1 \end{pmatrix} + \lambda \begin{pmatrix} 1 \\ -2 \\ 1 \end{pmatrix}$.

(a) Find the acute angle between the directions of AB and l. [4]

(b) Find the position vector of the point P on l such that $AP = BP$. [5]

9 The equation of a curve is $y = x^{-\frac{2}{3}} \ln x$ for $x > 0$. The curve has one stationary point.

(a) Find the exact coordinates of the stationary point. [5]

(b) Show that $\displaystyle\int_1^8 y \, \mathrm{d}x = 18 \ln 2 - 9$. [5]

10 The variables x and t satisfy the differential equation $\dfrac{\mathrm{d}x}{\mathrm{d}t} = x^2(1 + 2x)$, and $x = 1$ when $t = 0$.

Using partial fractions, solve the differential equation, obtaining an expression for t in terms of x. [11]