Cambridge Assessment International Education

Cambridge International AS & A Level

CANDIDATE NAME	
CENTRE NUMBER	
CANDIDATE NUMBER	

5794103083

MATHEMATICS **9709/31**

Paper 3 Pure Mathematics 3 **May/June 2022**

1 hour 50 minutes

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [ ].

This document has 4 pages.

© UCLES 2022

1 Solve the equation $2(3^{2x-1}) = 4^{x+1}$, giving your answer correct to 2 decimal places. [4]

2 **(a)** Expand $\left(2 - x^2\right)^{-2}$ in ascending powers of x, up to and including the term in x^4, simplifying the coefficients. [4]

(b) State the set of values of x for which the expansion is valid. [1]

3 Solve the equation $2\cot 2x + 3\cot x = 5$, for $0° < x < 180°$. [6]

4 The variables x and y satisfy the differential equation

$$\frac{\mathrm{d}y}{\mathrm{d}x} = \frac{xy}{1+x^2},$$

and $y = 2$ when $x = 0$.

Solve the differential equation, obtaining a simplified expression for y in terms of x. [7]

5 The polynomial $ax^3 - 10x^2 + bx + 8$, where a and b are constants, is denoted by $\mathrm{p}(x)$. It is given that $(x - 2)$ is a factor of both $\mathrm{p}(x)$ and $\mathrm{p}'(x)$.

(a) Find the values of a and b. [5]

(b) When a and b have these values, factorise $\mathrm{p}(x)$ completely. [3]

6 Let $I = \displaystyle\int_0^3 \frac{27}{\left(9+x^2\right)^2}\,\mathrm{d}x$.

(a) Using the substitution $x = 3\tan\theta$, show that $I = \displaystyle\int_0^{\frac{1}{4}\pi} \cos^2\theta\,\mathrm{d}\theta$. [4]

(b) Hence find the exact value of I. [4]

7 The complex number u is defined by $u = \dfrac{\sqrt{2} - a\sqrt{2}\mathrm{i}}{1 + 2\mathrm{i}}$, where a is a positive integer.

(a) Express u in terms of a, in the form $x + \mathrm{i}y$, where x and y are real and exact. [3]

It is now given that $a = 3$.

(b) Express u in the form $re^{\mathrm{i}\theta}$, where $r > 0$ and $-\pi < \theta \leqslant \pi$, giving the exact values of r and θ. [2]

(c) Using your answer to part **(b)**, find the two square roots of u. Give your answers in the form $re^{i\theta}$, where $r > 0$ and $-\pi < \theta \leqslant \pi$, giving the exact values of r and θ. [3]


8 The equation of a curve is $x^3 + y^3 + 2xy + 8 = 0$.

(a) Express $\frac{dy}{dx}$ in terms of x and y. [4]

The tangent to the curve at the point where $x = 0$ and the tangent at the point where $y = 0$ intersect at the acute angle α.

(b) Find the exact value of $\tan\alpha$. [5]

9




In the diagram, $OABCDEFG$ is a cuboid in which $OA = 2$ units, $OC = 4$ units and $OG = 2$ units. Unit vectors **i**, **j** and **k** are parallel to OA, OC and OG respectively. The point M is the midpoint of DF. The point N on AB is such that $AN = 3NB$.

(a) Express the vectors $\overrightarrow{OM}$ and $\overrightarrow{MN}$ in terms of **i**, **j** and **k**. [3]

(b) Find a vector equation for the line through M and N. [2]

(c) Show that the length of the perpendicular from O to the line through M and N is $\sqrt{\frac{53}{6}}$. [4]

10



The curve $y = x\sqrt{\sin x}$ has one stationary point in the interval $0 < x < \pi$, where $x = a$ (see diagram).

(a) Show that $\tan a = -\frac{1}{2}a$. [4]

(b) Verify by calculation that a lies between 2 and 2.5. [2]

(c) Show that if a sequence of values in the interval $0 < x < \pi$ given by the iterative formula $x_{n+1} = \pi - \tan^{-1}\left(\frac{1}{2}x_n\right)$ converges, then it converges to a, the root of the equation in part **(a)**. [2]

(d) Use the iterative formula given in part **(c)** to determine a correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

© UCLES 2022